

Savings
information
on back.

Making the most of your treatment

Learn about a prescription
medicine that can help relieve
moderate to severe chronic pain.

The use of oxycodone medications,
including Xtampza ER, may result
in serious, life-threatening, or fatal
respiratory depression even when
used as recommended.



*Proven pain
relief I can
count on*

INDICATIONS AND USAGE

Xtampza[®] ER (oxycodone) is:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain severe enough to require daily, around-the-clock, long-term treatment with an opioid when other pain treatments, such as non-opioid pain medicines or immediate-release opioid medicines, do not treat your pain well enough or you cannot tolerate them.
- A long-acting (extended-release) opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed by your healthcare provider, you are at risk for opioid addiction, abuse, and misuse that can lead to death.
- Not for use to treat pain that is not around-the-clock.

IMPORTANT SAFETY INFORMATION ABOUT XTAMPZA ER

- **Get emergency help or call 911 right away if you take too much Xtampza ER (overdose).** When you first start taking Xtampza ER, when your dose is changed, or if you take too much (overdose), serious life-threatening breathing problems that can lead to death may occur. Talk to your healthcare provider about naloxone, a medicine for the emergency treatment of an overdose.
- Taking Xtampza ER with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.
- Never give anyone else your Xtampza ER. They could die from taking it. Selling or giving away Xtampza ER is against the law.
- Store Xtampza ER securely, out of sight and reach of children, and in a location not accessible by others, including visitors to the home

Please see Important Safety Information and full Prescribing Information, including Boxed Warning on Addiction, Abuse, and Misuse and other serious risks accompanying this piece, or at XtampzaER.com/PI

What is chronic pain?

Everyone experiences pain from time to time. But chronic pain doesn't go away. Healthcare providers define it as pain that lasts longer than 3 months. Chronic pain could be caused by a health condition or injury, but sometimes there's not an obvious cause.

If you are one of the millions of Americans suffering from chronic pain, treatments such as Xtampza ER may be able to help you manage moderate to severe chronic pain.

What is Xtampza ER (oxycodone)?

Xtampza ER (pronounced ex tamp' zah ee ar) is

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain severe enough to require daily, around-the-clock, long-term treatment with an opioid when other pain treatments, such as non-opioid pain medicines or immediate-release opioid medicines, do not treat your pain well enough or you cannot tolerate them.
- A long-acting (extended-release) opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed by your healthcare provider, you are at risk for opioid addiction, abuse, and misuse that can lead to death.
- Not for use to treat pain that is not around-the-clock.

Keep in mind...

- Store Xtampza ER in a safe place, out of sight of others, and secured in a locked cabinet, drawer, or safe
- Never give anyone else this prescription medicine. It can be life-threatening. Selling or giving away Xtampza ER is against the law.

How is Xtampza® ER (oxycodone) different?


Xtampza ER is a form of oxycodone designed with extended-release technology. When taking Xtampza ER, here are some of the things you can expect:



Chronic pain relief that's released slowly over time, unlike short-acting pain medicines that may work quickly, but don't last as long.



Dosed every 12 hours, so you only take it twice a day. Xtampza ER should be taken at the same time every day and **MUST be taken with food.**



Chewing or crushing some pain medications can result in a dangerous spike of medication levels in your blood.

Xtampza ER offers the ability to open the capsule and sprinkle the medicine if you dislike or have trouble swallowing capsules without resulting in a dangerous spike in medication.

The use of any opioid may result in serious, life-threatening, or fatal respiratory depression, even when used as recommended.

Although Xtampza ER is formulated to make manipulation more difficult, it cannot entirely prevent abuse; abuse of Xtampza ER by injection and via the oral and nasal routes is still possible.

How to take Xtampza ER

You'll take Xtampza ER dose twice a day—every 12 hours—at the same times, and you **MUST take it with food.** You should take each dose with approximately the same amount of food each time to help the medicine work properly to relieve pain.

It helps to establish a pattern to help you remember to take it. For example, you can take it at 7 am with breakfast and 7 pm with dinner each day.

- **If Xtampza ER doesn't control your pain, talk to your healthcare provider about adjusting your treatment plan. Never change your dose or stop taking Xtampza ER without talking to your healthcare provider first.**

Please see Important Safety Information and full Prescribing Information, including Boxed Warning on Addiction, Abuse, and Misuse and other serious risks accompanying this piece, or at XtampzaER.com/PI

Xtampza® ER
(oxycodone) EXTENDED-RELEASE CAPSULES 

You have 4 options for taking your medicine:



The whole capsule by mouth



Sprinkle on soft foods

(for example: applesauce, pudding, yogurt, ice cream, or jam)



Sprinkle into a cup and then swallow



Through a gastrostomy/nasogastric tube

Xtampza® ER MUST be taken with food.

Please see the Instructions for Use in the enclosed Prescribing Information for more details, and talk to your healthcare provider.

How should you store your prescription?

Store your prescription at room temperature, away from light. It's important to keep it in a safe place, out of sight and secured in a locked cabinet, drawer, or safe.

While taking Xtampza® ER (oxycodone), DO NOT:

- Drive or operate heavy machinery until you know how Xtampza ER affects you. Xtampza ER can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or take over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with Xtampza ER may cause you to overdose and die.

Do not take Xtampza ER if you have:

- Severe asthma
- Trouble breathing or other lung problems
- A bowel blockage
- Narrowing of the stomach or intestines

Talking with your healthcare provider

- Before you start taking Xtampza ER, tell your healthcare provider if you have a history of:
 - head injury
 - seizures
 - liver problems
 - kidney problems
 - thyroid problems
 - problems urinating
 - pancreas or gall bladder problems
 - abuse of street or prescription drugs, alcohol addiction, opioid overdose, or mental health problems
- Tell your healthcare provider if you're:
 - **pregnant or planning to become pregnant**
 - **breastfeeding**
 - living in a household where there are small children or someone who has abused street or prescription drugs.
 - taking prescription or over-the-counter medicines, vitamins, or herbal supplements.

Taking Xtampza ER with certain other medicines can cause serious side effects that could lead to death.

Did you know?

Xtampza ER is the only extended-release oxycodone in a capsule that can be opened and sprinkled. With other extended-release pain medicines, chewing or crushing them can disrupt the dosage, destroy the extended-release properties, and increase the risk of side effects for some people.

Please see Important Safety Information and full Prescribing Information, including Boxed Warning on Addiction, Abuse, and Misuse and other serious risks accompanying this piece, or at XtampzaER.com/PI



Possible side effects of Xtampza® ER to watch for

- Constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, and abdominal pain.
- Call your healthcare provider if you have any of these symptoms and they are severe.

Get emergency medical help if you have any of these symptoms:

- trouble breathing
- shortness of breath
- fast heartbeat
- chest pain
- swelling of your face, tongue, or throat
- extreme drowsiness
- light-headedness when changing positions
- feeling faint
- feeling agitated
- high body temperature
- trouble walking
- stiff muscles
- mental changes such as confusion

These are not all the possible side effects of Xtampza ER. Call your healthcare provider for medical advice about side effects.



The side effects seen with Xtampza ER are similar to those seen with other oxycodone medications.

There are no head-to-head studies comparing the safety and efficacy of Xtampza ER to other oxycodone medications.

Important information to keep in mind

- **Get emergency help or call 911 right away if you take too much Xtampza ER (overdose).** When you first start taking Xtampza ER, when your dose is changed, or if you take too much (overdose), serious life-threatening breathing problems that can lead to death may occur. Talk to your healthcare provider about naloxone, a medicine for the emergency treatment of an overdose.
- Taking Xtampza ER with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.
- Never give anyone else your Xtampza ER. They could die from taking it. Store Xtampza ER away from children and in a safe place to prevent stealing or abuse. Selling or giving away Xtampza ER is against the law.
- Store Xtampza ER securely, out of sight and reach of children, and in a location not accessible by others, including visitors to the home.

Are you getting the most out of your treatment?



Work together with your healthcare provider.

Talk to your healthcare provider about how your treatment is going. Never change or stop taking your dose without talking to your healthcare provider first.



Stick to the plan.

Always follow your healthcare provider's instructions. Take Xtampza ER with the same amount of food to make sure the medicine is consistently absorbed.



Find what works for you.

If you'd prefer to take your medicine a certain way (if you don't like swallowing pills, for example), talk with your healthcare provider about your options.



Stay on schedule.

Remember to take your prescribed dose every 12 hours, at the same times every day. If you miss a dose, take your next dose at your usual time.

Did you know?

It's important to learn all you can about living with chronic pain. These organizations can help you find support groups in your area and also provide other helpful information.

American Academy of Pain Medicine*

<https://painmed.org/organizations-dedicated-to-people-with-pain/>

American Chronic Pain Association*

<https://www.acpanow.com/>

National Institute of Health's Medline Plus

<https://medlineplus.gov/backpain.html>

*Collegium provides financial support to these organizations
Collegium Pharmaceutical, Inc. is not responsible for and does not create, control, fund, or influence any of the content found on any third-party website

Please see Important Safety Information and full Prescribing Information, including Boxed Warning on Addiction, Abuse, and Misuse and other serious risks accompanying this piece, or at XtampzaER.com/PI

Xtampza[®] ER
(oxycodone) EXTENDED-RELEASE CAPSULES 

Responsible opioid usage

The opioid crisis is a major problem facing communities throughout our country. As part of this crisis, prescription opioids have been stolen, misused and abused.

You've been prescribed Xtampza® ER (oxycodone), an abuse-deterrent opioid that was developed with these issues in mind, and it's important that you understand how these issues can impact you. Ask your doctor for more information about how Xtampza ER can help.

Another way to address the opioid epidemic is by responsibly storing and disposing of your medication. Make sure you are:

- Taking Xtampza ER exactly as your doctor has prescribed
- Storing your medications safely and securely (consider placing all medication in a locked box or cabinet)
- Discarding unused medications properly and not saving them

Proper disposal procedures include:

Take-back programs

- Visit takebackday.dea.gov for National Prescription Drug Take-Back events hosted by the DEA in communities nationwide.
- Your local law enforcement agencies may also sponsor medicine take-back programs. For more information, contact your city or county government.

Authorized collectors

Retail pharmacies, hospital or clinic pharmacies, and law enforcement locations can be authorized collection sites. To find authorized collectors near you:

- Visit <https://apps.DEAdiversion.usdoj.gov/pubdispsearch/>
- Call the DEA's Registration Call Center at 1-800-882-9539

If these options are not available to you, you can flush your unused Xtampza ER capsules down the toilet.

IMPORTANT SAFETY INFORMATION ABOUT XTAMPZA® ER (oxycodone)

WARNING: ADDICTION, ABUSE, AND MISUSE; RISK EVALUATION AND MITIGATION STRATEGY (REMS); LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; NEONATAL OPIOID WITHDRAWAL SYNDROME; CYTOCHROME P450 3A4 INTERACTION; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

Addiction, Abuse, and Misuse

Xtampza ER exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Assess each patient's risk prior to prescribing Xtampza ER and monitor all patients regularly for the development of these behaviors or conditions.

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)

To ensure that the benefits of opioid analgesics outweigh the risks of addiction, abuse, and misuse, the Food and Drug Administration (FDA) has required a REMS for these products. Under the requirements of the REMS, drug companies with approved opioid analgesic products must make REMS-compliant education programs available to healthcare providers. Healthcare providers are strongly encouraged to

- complete a REMS-compliant education program,
- counsel patients and/or their caregivers, with every prescription, on safe use, serious risks, storage, and disposal of these products,
- emphasize to patients and their caregivers the importance of reading the Medication Guide every time it is provided by their pharmacist, and
- consider other tools to improve patient, household, and community safety.

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of Xtampza ER. Monitor for respiratory depression, especially during initiation of Xtampza ER or following a dose increase.

Accidental Ingestion

Accidental ingestion of even one dose of Xtampza ER, especially by children, can result in a fatal overdose of oxycodone.

Neonatal Opioid Withdrawal Syndrome

Prolonged use of Xtampza ER during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available.

Cytochrome P450 3A4 Interaction

The concomitant use of Xtampza ER with all cytochrome P450 3A4 inhibitors may result in an increase in oxycodone plasma concentrations, which could increase or prolong adverse drug effects and may cause potentially fatal respiratory depression. In addition, discontinuation of a concomitantly used cytochrome P450 3A4 inducer may result in an increase in oxycodone plasma concentration. Monitor patients receiving Xtampza ER and any CYP3A4 inhibitor or inducer.

Risks From Concomitant Use With Benzodiazepines or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death.

- Reserve concomitant prescribing of Xtampza ER and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.
- Limit dosages and durations to the minimum required.
- Follow patients for signs and symptoms of respiratory depression and sedation.

Please see Important Safety Information and full Prescribing Information, including Boxed Warning on Addiction, Abuse, and Misuse and other serious risks accompanying this piece, or at XtampzaER.com/PI



IMPORTANT SAFETY INFORMATION ABOUT XTAMPZA® ER continued

Important Information about Xtampza ER

- **Get emergency help or call 911 right away if you take too much Xtampza ER (overdose).** When you first start taking Xtampza ER, when your dose is changed, or if you take too much (overdose), serious life-threatening breathing problems that can lead to death may occur. Talk to your healthcare provider about naloxone, a medicine for the emergency treatment of an overdose.
- Taking Xtampza ER with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.
- Never give anyone else your Xtampza ER. They could die from taking it. Selling or giving away Xtampza ER is against the law.
- Store Xtampza ER securely, out of sight and reach of children, and in a location not accessible by others, including visitors to the home.

Do not take Xtampza ER if you have:

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage, or have narrowing of the stomach or intestines.

Before taking Xtampza ER, tell your healthcare provider if you have a history of:

- head injury, seizures
- liver, kidney, thyroid problems
- problems urinating
- pancreas or gallbladder problems
- abuse of street or prescription drugs, alcohol addiction, opioid overdose, or mental health problems

Tell your healthcare provider if you are:

- **pregnant or planning to become pregnant.** Prolonged use of Xtampza ER during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **breastfeeding.** Not recommended during treatment with Xtampza ER. It may harm your baby.
- living in a household where there are small children or someone who has abused street or prescription drugs.
- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking Xtampza ER with certain other medicines can cause serious side effects that could lead to death.

When taking Xtampza ER:

- Do not change your dose. Take Xtampza ER exactly as prescribed by your healthcare provider. Use the lowest dose possible for the shortest time needed.
- Take your prescribed dose every 12 hours, at the same time every day. Do not take more than your prescribed dose. If you miss a dose, take your next dose at your usual time.

- If you cannot swallow Xtampza ER capsules, see the detailed Instructions for Use in the Medication Guide.
- Always take Xtampza ER capsules with approximately the same amount of food to ensure enough medicine is absorbed.
- Swallow Xtampza ER whole. Do not snort, or inject Xtampza ER because this may cause you to overdose and die.
- The contents of the Xtampza ER capsules may be sprinkled on soft food, sprinkled into a cup and then put directly into the mouth, or given through a nasogastric or gastrostomy tube.
- **Call your healthcare provider if the dose you are taking does not control your pain.**
- **Do not stop taking Xtampza ER without talking to your healthcare provider.**
- Dispose of expired, unwanted or unused Xtampza ER by promptly flushing down the toilet, if a drug take-back option is not readily available. Visit www.fda.gov/drugdisposal for additional information on disposal of unused medicines.

While taking Xtampza ER, DO NOT:

- Drive or operate heavy machinery, until you know how Xtampza ER affects you. Xtampza ER can make you sleepy, dizzy, or light-headed.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with Xtampza ER may cause you to overdose and die.

The possible side effects of Xtampza ER are:

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, and abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe.

Get emergency medical help or call 911 right away if you have:

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion.

These are not all the possible side effects of Xtampza ER. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. **For more information, go to dailymed.nlm.nih.gov.**

Please see Important Safety Information and full Prescribing Information, including Boxed Warning on Addiction, Abuse, and Misuse and other serious risks accompanying this piece, or at XtampzaER.com/PI. Speak to your healthcare provider if you have questions about Xtampza ER.



Start saving today with the Xtampza® ER Co-pay Card



The image shows a green co-pay card. On the left is a small graphic of a pill box with an 'X' on it. In the top right corner is the Xtampza ER logo, which includes the text 'Xtampza ER' and '(oxycodone) EXTENDED-RELEASE CAPSULES II'. The main text on the card reads: 'Pay as little as \$35* for each prescription'.

Eligible patients may pay as little as \$35* for each prescription.



Visit XtampzaER.com for additional information and resources.

CO-PAY SAVINGS* available for eligible commercially insured patients.

*Please see Program Terms, Conditions, and Eligibility Criteria at www.XtampzaER.com or scan the QR code. Maximum savings limit applies; patient out-of-pocket expense may vary.



For questions about Xtampza ER, please call 888-884-2655. Representatives available Monday through Friday, 9AM – 5PM EST.

Program Terms, Conditions, and Eligibility Criteria

Offer is valid for commercially insured patients only. Most eligible commercially insured patients will pay as little as \$35 (maximum benefit amount of \$125) for each fill of Xtampza® ER (oxycodone) capsules. Any remaining out-of-pocket expense will be the patient's responsibility. This offer is valid in the United States. Some restrictions may apply. Offer not valid for prescriptions reimbursed under Medicaid, a Medicare drug benefit plan, Tricare or other federal or state health programs (such as medical assistance programs). Cash Discount Cards and other non-insurance plans are not valid as primary under this offer. If the patient is eligible for drug benefits under any such program, the patient cannot use this offer. By using this offer, the patient certifies that he or she will comply with any terms of his or her health insurance contract requiring notification to his or her payer of the existence and/or value of this offer. This offer is not valid for cash paying patients. Offer not valid for patients under 18 years of age. **Program is not valid in Massachusetts or any other state or jurisdiction where prohibited or restricted by law.**

Patient Instructions: To redeem this offer, you must have a valid prescription for Xtampza® ER and follow the dosage instructions provided by your healthcare provider with a valid Prescriber ID# (e.g. NPI or DEA). This offer may not be redeemed for cash. Patients with questions about the Xtampza® ER savings offer should call **1-844-247-4753**.

This offer is not transferable and is limited to one offer per person. Not valid if reproduced. Void where prohibited by law. Program managed by ConnectiveRx on behalf of Collegium Pharmaceutical. The parties reserve the right to rescind, revoke or amend this offer without notice at any time.



See Important Safety Information and full Prescribing Information, including Boxed Warning on Addiction, Abuse, and Misuse and other serious risks accompanying this piece, or at [Xtampza ER.com/PI](http://XtampzaER.com/PI). Speak to your healthcare provider if you have questions about Xtampza ER.

XTAMPZA, Collegium Pharmaceutical, and the Collegium Pharmaceutical logo are registered trademarks of Collegium Pharmaceutical, Inc.

© 2023 Collegium Pharmaceutical, Inc. All rights reserved. PP-XT-US-1110 07/23